

7B TestOfHyp

Contents

A Test of Hypothesis

A.1 Example	
A.2 figure	
A.3 figure	
A.4 Standardize the statistic	
A.5 fig	
A.6 Critical Value	
A.7 fig	
A.8 Under Alternative	
A.9 fig	
A.10 Type I and Type II Errors	
A.11 P-value vs α	
A.12 Test of Hypothesis - upper-tailed alternative	
A.13 Test of Hypothesis - lower-tailed alternative	
A.14 Test of Hypothesis - two-tailed alternative	

A Test of Hypothesis

[ToC]

A.1 Example

- Suppose we want to test two hypothesis

$$H_0 :$$

$$p = .5$$


$$H_A :$$

$$p = .6$$


- How can we test these hypothesis?

A.2 figure

n=50: If p=.5




If p=.6




A.3 figure


$n=100$: If $p=.5$




If $p=.6$



n=500: If p=.5



If p=.6



A.4 Standardize the statistic


- Instead of looking at X , let's look at the z-score of $\hat{p} = X/n$.

$$Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$$


$$\frac{(.6 - .5)}{\sqrt{(.5)(.5)/50}} = 1.414 \quad \frac{(.6 - .5)}{\sqrt{(.5)(.5)/100}} = 2 \quad \frac{(.6 - .5)}{\sqrt{(.5)(.5)/500}} = 4.47$$

A.5 fig


n=50: If p=.5




If p=.6




$n=100$: If $p=.5$




If $p=.6$



n=500: If p=.5



If p=.6




A.6 Critical Value


- Now that the distribution is fixed under H_0 , we can fix the critical value (blue line) as well:

A.7 fig


$n=50$: If $p=.5$




If $p=.6$




n=100: If p=.5




If p=.6



n=500: If p=.5




If p=.6




A.8 Under Alternative

- What if the friend was cheating by different amount?


n=50: If p=.5




If p=.55




n=50: If p=.5




If p=.6



n=50: If p=.5




If p=.65



A.9 fig


$n=100$: If $p=.5$




If $p=.55$




n=100: If p=.5




If p=.6



n=100: If p=.5



If p=.65




A.10 Type I and Type II Errors

- With significance level α of our choice,

Prob of ...	don't reject H_0	reject H_0
H_0 is true	$1 - \alpha$	(α)
H_A is true	(β)	Power = $1 - \beta$

- $\alpha = P(\text{ type I error }) = P(\text{ false positive})$
- $\beta = P(\text{ type II error }) = P(\text{ false negative})$

A.11 P-value vs α



- p-value is the probability of getting the observed value of test statistics z or 'worse' when H_0 is true.

A.12 Test of Hypothesis - upper-tailed alternative

1. Set up the null and alternative hypothesis

$$H_0 : \quad p = .5$$

$$H_A : \quad p > .5$$

2. Calculate the test statistic



$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

3. Get p-value

4. Conclude

If p-value is greater than α

If p-value is not greater than α



A.13 Test of Hypothesis - lower-tailed alternative

1. Set up the null and alternative hypothesis

$$H_0 : \quad p = .5$$

$$H_A : \quad p < .5$$

2. Calculate the test statistic

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

3. Get p-value

4. Conclude

If p-value is greater than α

If p-value is less than α

A.14 Test of Hypothesis - two-tailed alternative

1. Set up the null and alternative hypothesis

$$H_0 : p = .5$$

$$H_A : p \neq .5$$

2. Calculate the test statistic

$$z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$$

3. Get p-value

4. Conclude

If p-value is greater than α

If p-value is less than α

